

Local action of the symmetric group and the twisted Yang-Baxter relation

Alexander Odesskii

Introduction

The Yang-Baxter relation plays a central role in the Quantum Field Theory. The usual R -matrix describes the scattering of two "particles" (Fig.1). Here u and v are some characteristics of the particles which are called the spectral parameters. R -matrix ($R_{ij}^{kl}(u, v)$) is a meromorphic function on the space of spectral parameters. Now suppose that the spectral parameters change after scattering (Fig.2). New spectral parameters $\varphi(u, v)$, $\psi(u, v)$ are some meromorphic functions in the variables u and v . The factorization conditions impose a strong requirement on these functions (Fig.3). One can see from Fig.3 that

$$\varphi(u, \varphi(v, w)) = \varphi(\varphi(u, v), \varphi(\psi(u, v), w))$$

$$\varphi(\psi(u, \varphi(v, w)), \psi(v, w)) = \psi(\varphi(u, v), \varphi(\psi(u, v), w))$$

$$\psi(\psi(u, v), w) = \psi(\psi(u, \varphi(v, w)), \psi(v, w))$$

If we have the solution of these equations, then we can define the analogous Yang-Baxter relation, L -operators, Zamolodchikov algebra and so on. In §1 we develop the mathematical formalism for φ and ψ and give nontrivial examples. In §2 we introduce the twisted Yang-Baxter relation and corresponding algebraic structures. The most interesting examples of φ and ψ appear in the theory of factorization of matrix polynomials (§3) and matrix θ -functions (§4). In §5 we discuss the generalized star-triangle relation and corresponding algebraic structures. In "physical" language it is the case when our particles are not elementary but are multiplets of m elementary particles.




Figure 1




Figure 2

The examples of twisted R -matrices as well as the solutions of generalized star-triangle relation were found in [4] as intertwiners of cyclic representations and its tensor products of the algebra of monodromy matrices of the six-vertex model at root of unity [3]. These solutions are natural generalizations of the one from chiral

Potts model [1,2,3]. One can obtain other solutions of the twisted Yang-Baxter and star-triangle relations by calculating the intertwiners of the representations of the algebras of monodromy matrices at root of unity for other trigonometric and elliptic R -matrices.




Figure 3

§1. Twisted transposition

Let U be a complex manifold, $\mu : U \times U \rightarrow U \times U$ be a birational automorphism of $U \times U$. We will use the notations: $\mu(u, v) = (\varphi(u, v), \psi(u, v))$ where $u, v \in U$. Here φ and ψ are meromorphic functions from $U \times U$ to U .

Let us introduce the following birational automorphisms of $U \times U \times U$: $\sigma_1 = \mu \times \text{id}$ and $\sigma_2 = \text{id} \times \mu$. We have: $\sigma_1(u, v, w) = (\varphi(u, v), \psi(u, v), w)$ and $\sigma_2(u, v, w) = (u, \varphi(v, w), \psi(v, w))$.

Definition We call a map μ a twisted transposition if the automorphisms σ_1 and σ_2 satisfy the following relations:

$$\sigma_1^2 = \sigma_2^2 = \text{id}, \quad \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \quad (1)$$

If μ is a twisted transposition, then for each $N \in \mathbb{N}$ we have the birational action of the symmetric group S_N on the manifold U^N such that the transposition $(i, i+1)$ acts by automorphism $\sigma_i = \text{id}^{i-1} \times \mu \times \text{id}^{N-i-1}$. So we have $\sigma_i(u_1, \dots, u_N) = (u_1, \dots, \varphi(u_i, u_{i+1}), \psi(u_i, u_{i+1}), \dots, u_N)$. It is clear that the relations (1) are equivalent to the defining relations in the group S_N : $\sigma_i^2 = e, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \sigma_i \sigma_j = \sigma_j \sigma_i$ for $|i - j| > 1$.

It is easy to check that the relations (1) are equal to the following functional equations for φ and ψ :

$$\begin{aligned} \varphi(\varphi(u, v), \psi(u, v)) &= u, \quad \psi(\varphi(u, v), \psi(u, v)) = v \\ \varphi(u, \varphi(v, w)) &= \varphi(\varphi(u, v), \varphi(\psi(u, v), w)) \\ \varphi(\psi(u, \varphi(v, w)), \psi(v, w)) &= \psi(\varphi(u, v), \varphi(\psi(u, v), w)) \\ \psi(\psi(u, v), w) &= \psi(\psi(u, \varphi(v, w)), \psi(v, w)) \end{aligned} \quad (2)$$

Remarks 1. From (2) it follows that for each N the functions $\varphi(u_1, \varphi(u_2, \dots, \varphi(u_N, w) \dots))$ and $\psi(\dots(\psi(w, u_1), u_2) \dots, u_N)$ are invariant with respect to the action of the group S_N on the variables $u_1, \dots, u_N$.

2. Informally one can consider μ as an infinite dimensional R -matrix in the space of functions. Namely, if we consider the space of meromorphic functions $\{f : U \times U \rightarrow \mathbb{C}\}$ as an "extended tensor square" of the space of meromorphic functions $\{f : U \rightarrow \mathbb{C}\}$, then the linear operator $R_\mu : f \rightarrow f \circ \mu$ (that is $R_\mu f(u, v) = f(\mu(u, v))$) satisfies the usual Yang-Baxter relation.

Examples 1. Let $q, q^{-1} : U \rightarrow U$ be birational automorphisms such that $qq^{-1} = q^{-1}q = \text{id}$. Then

$$\mu(u, v) = (q(v), q^{-1}(u))$$

is a twisted transposition.

2. Let $U = \mathbb{C}$, then the following formula gives a twisted transposition:

$$\mu(u, v) = (1 - u + uv, \frac{uv}{1 - u + uv})$$

3. Let U be a finite dimensional associative algebra with unity $1 \in U$ and such that generic element $u \in U$ is invertible, for example $U = \text{Mat}_m$. Then the following formula gives a twisted transposition:

$$\mu(u, v) = (1 - u + uv, (1 - u + uv)^{-1}uv)$$

For further examples see §**3,4**.

§2. Twisted Yang-Baxter relation and corresponding algebraic structures

Let V be a n -dimensional vector space. For each $u \in U$ we denote by $V(u)$ a vector space canonically isomorphic to V . Let R be a meromorphic function from $U \times U$ to $End(V \otimes V)$. We will consider $R(u, v)$ as a linear operator

$$R(u, v) : V(u) \otimes V(v) \rightarrow V(\varphi(u, v)) \otimes V(\psi(u, v))$$

Definition We call R a twisted R -matrix (with respect to the twisted transposition μ) if it satisfies the following properties:

1. The composition

$$V(u) \otimes V(v) \rightarrow V(\varphi(u, v)) \otimes V(\psi(u, v)) \rightarrow V(u) \otimes V(v)$$

is equal to the identity, that is $R(\varphi(u, v), \psi(u, v))R(u, v) = 1$.

2. The following diagram is commutative:

$$\begin{array}{ccc}
 V(\varphi(u, v)) \otimes V(\psi(u, v)) \otimes V(w) & \xrightarrow{1 \otimes R(\psi(u, v), w)} & V(\varphi(u, v)) \otimes V(\varphi(\psi(u, v), w)) \otimes V(\psi(\psi(u, v), w)) \\
 \uparrow R(u, v) \otimes 1 & & \downarrow R(\varphi(u, v), \varphi(\psi(u, v), w)) \otimes 1 \\
 V(u) \otimes V(v) \otimes V(w) & & \tilde{V} \\
 \downarrow 1 \otimes R(v, w) & & \uparrow 1 \otimes R(\psi(u, \varphi(v, w)), \psi(v, w)) \\
 V(u) \otimes V(\varphi(v, w)) \otimes V(\psi(v, w)) & \xrightarrow{R(u, \varphi(v, w)) \otimes 1} & V(\varphi(u, \varphi(v, w))) \otimes V(\psi(u, \varphi(v, w))) \otimes V(\psi(v, w))
 \end{array}$$

Here $\tilde{V} = V(\varphi(u, \varphi(v, w))) \otimes V(\psi(\varphi(u, v), \varphi(\psi(u, v), w))) \otimes V(\psi(\psi(u, v), w))$.

In other words,

$$\begin{aligned}
 R^{12}(\varphi(u, v), \varphi(\psi(u, v), w)) R^{23}(\psi(u, v), w) R^{12}(u, v) = \\
 R^{23}(\psi(u, \varphi(v, w)), \psi(v, w)) R^{12}(u, \varphi(v, w)) R^{23}(v, w)
 \end{aligned} \tag{3}$$

Here $R^{12} = R \otimes 1$ and $R^{23} = 1 \otimes R$ are linear operators in $V \otimes V \otimes V$.

We call (3) twisted Yang-Baxter relation.

Let $\{x_i, i = 1 \dots, n\}$ be a basis of the linear space V , $\{x_i(u)\}$ be the corresponding basis of the linear space $V(u)$. It is clear that the following two linear operators are twisted R -matrices for each μ :

$$x_i(u) \otimes x_j(v) \rightarrow x_i(\varphi(u, v)) \otimes x_j(\psi(u, v))$$

$$x_i(u) \otimes x_j(v) \rightarrow x_j(\varphi(u, v)) \otimes x_i(\psi(u, v))$$

Let $R_{ij}^{\alpha\beta}(u, v)$ be a matrix element of twisted R -matrix R , which means

$$R(u, v) : x_i(u) \otimes x_j(v) \rightarrow R_{ij}^{\alpha\beta}(u, v) x_\beta(\varphi(u, v)) \otimes x_\alpha(\psi(u, v))$$

We sum over repeating indices.

Definition We call the twisted Zamolodchikov algebra associated with R an associative algebra Z_R with generators $\{x_i(u); i = 1 \dots, n, u \in U\}$ and defining relations

$$x_i(u) x_j(v) = R_{ij}^{\alpha\beta}(u, v) x_\beta(\varphi(u, v)) x_\alpha(\psi(u, v))$$

It is clear that the twisted Zamolodchikov algebra satisfies the usual property: for generic $u_1, \dots, u_N \in U$ the elements $\{x_{i_1}(u_1) \dots x_{i_N}(u_N); 1 \leq i_1, \dots, i_N \leq n\}$ are linear independent.

Let W be a vector space, L be a meromorphic function $L : U \rightarrow \text{End}(V \otimes W)$. We will consider $L(u)$ as a linear operator $L(u) : V(u) \otimes W \rightarrow W \otimes V(u)$.

Definition We call L an L -operator for twisted R -matrix R if the following diagram is commutative:

$$\begin{array}{ccc}
V(u) \otimes W \otimes V(v) & \xrightarrow{L(u) \otimes 1} & W \otimes V(u) \otimes V(v) \\
\uparrow 1 \otimes L(v) & & \downarrow 1 \otimes R(u, v) \\
V(u) \otimes V(v) \otimes W & & W \otimes V(\varphi(u, v)) \otimes V(\psi(u, v)) \\
\downarrow R(u, v) \otimes 1 & & \uparrow L(\varphi(u, v)) \otimes 1 \\
V(\varphi(u, v)) \otimes V(\psi(u, v)) \otimes W & \xrightarrow{1 \otimes L(\psi(u, v))} & V(\varphi(u, v)) \otimes W \otimes V(\psi(u, v))
\end{array}$$

Which is

$$R(u, v) L^1(u) L^2(v) = L^1(\varphi(u, v)) L^2(\psi(u, v)) R(u, v) \quad (4)$$

where $L^1 = L \otimes 1$ and $L^2 = 1 \otimes L$.

Let $\{L_i^j(u); i, j = 1, \dots, n\}$ be the elements in $\text{End}(W)$ such that $L(u) : x_i(u) \otimes W \rightarrow (L_i^\alpha(u)W) \otimes x_\alpha(u)$. We can consider $L(u)$ as a $n \times n$ matrix with matrix elements $(L_i^j(u))$.

Definition The algebra of monodromy matrices M_R for the twisted R -matrix R is an associative algebra with generators $\{L_i^j(u); i, j = 1, \dots, n, u \in U\}$ and defining relations (4).

It is clear that representations of the algebra M_R in a vector space W coincide with L -operators $L(u) \in \text{End}(V \otimes W)$.

One can introduce a coproduct Δ in the algebra M_R by the usual way:

$$\Delta L_i^j(u) = L_i^\alpha(u) \otimes L_\alpha^j(u)$$

In the language of L -operators, if we have two L -operators $L_1(u) : V(u) \otimes W_1 \rightarrow W_1 \otimes V(u)$ and $L_2(u) : V(u) \otimes W_2 \rightarrow W_2 \otimes V(u)$, then we can construct an L -operator in the space $W_1 \otimes W_2$ as the composition:

$$V(u) \otimes W_1 \otimes W_2 \rightarrow W_1 \otimes V(u) \otimes W_2 \rightarrow W_1 \otimes W_2 \otimes V(u)$$

Definition Q -operator is the element of the algebra M_R given by the formula:

$$Q(u) = \text{tr} L(u) = L_i^i(u)$$

From relations (4) it follows that Q -operators satisfy the following relations:

$$Q(u)Q(v) = Q(\varphi(u, v))Q(\psi(u, v))$$

Remark One can also define the dynamical twisted Yang-Baxter relation.

§3. Local action of the symmetric group and factorization of matrix polynomials

For the general theory of matrix polynomials and factorization see [7]. For our purposes we state the results, which may be well known to the experts.

We denote by $S(a)$ the set of eigenvalues of a matrix $a \in \text{Mat}_m$. More generally, we denote by $S(a_1, \dots, a_d)$, $a_1, \dots, a_d \in \text{Mat}_m$, the set of roots of the polynomial $f(t) = \det(t^d - a_1 t^{d-1} + \dots + (-1)^d a_d)$. We will consider polynomials with generic coefficients only, so $\#S(a_1, \dots, a_d) = md$.

Proposition 1. *Let*

$$t^d - a_1 t^{d-1} + \dots + (-1)^d a_d = (t - b_1) \dots (t - b_d) \quad (5)$$

for generic matrices $a_1, \dots, a_d \in \text{Mat}_m$, then $S(b_i) \cap S(b_j) = \emptyset$ for $i \neq j$ and $S(b_1) \cup \dots \cup S(b_d) = S(a_1, \dots, a_d)$. For each decomposition $S(a_1, \dots, a_d) = A_1 \cup \dots \cup A_d$, such that $\#A_i = m$, $A_i \cap A_j = \emptyset$ ($i \neq j$) there exists a unique factorization (5) with $S(b_i) = A_i$.

Proof The first statement follows from the equation $\det(t^d - a_1 t^{d-1} + \dots + (-1)^d a_d) = \det(t - b_1) \dots \det(t - b_d)$.

On the other hand, if we know eigenvalues of $b_1, \dots, b_d$ then we can calculate eigenvectors of them. For $\lambda \in S(b_d)$ the corresponding eigenvector is a vector v_λ , such that $(\lambda^d - a_1 \lambda^{d-1} + \dots + (-1)^d a_d) v_\lambda = 0$. If we know all eigenvectors of b_d , then we can calculate all eigenvectors of b_{d-1} similarly and so on. This implies the uniqueness. By our construction of $b_1, \dots, b_d$ the determinants of the matrix polynomials in the right hand side and the left hand side of (5) have the same set of roots. Moreover, for each root λ the operators represented by these matrix polynomials have the same kernel if we set $t = \lambda$. It implies that these polynomials are equal.

Proposition 2. *Let $a_1, a_2 \in \text{Mat}_m$ be generic matrices. Then there exists a unique pair of matrices $b_1, b_2 \in \text{Mat}_m$ such that $(t - a_1)(t - a_2) = (t - b_1)(t - b_2)$ and $S(b_1) = S(a_2), S(b_2) = S(a_1)$. We have $b_1 = a_1 + \Lambda^{-1}, b_2 = a_2 - \Lambda^{-1}$ where $a_2 \Lambda - \Lambda a_1 = 1$.*

Proof If $S(b_2) \cap S(a_2) \neq \emptyset$, then $\det(a_2 - b_2) = 0$, because a_2 and b_2 have a common eigenvector. Otherwise, we can put $\Lambda = (a_2 - b_2)^{-1}$.

Let $U = \text{Mat}_m$. From the propositions 1 and 2 it follows that the formula $\mu(a_1, a_2) = (b_1, b_2)$ gives a twisted transposition, where $b_1 + b_2 = a_1 + a_2$, $b_1 b_2 = a_1 a_2$, $S(b_1) = S(a_2), S(b_2) = S(a_1)$. We have $\mu(a_1, a_2) = (a_1 + \Lambda^{-1}, a_2 - \Lambda^{-1})$, where Λ is the solution of the linear matrix equation $a_2 \Lambda - \Lambda a_1 = 1$.

Let $\overline{U} = \overline{\text{Mat}}_m^N$ be the set of $m \times m$ matrices with different eigenvalues and fixed order of eigenvalues. The proposition 1 gives an action of the symmetric group S_{mN} on the space $\overline{U}^N$ by birational automorphisms. By definition, for $\sigma \in S_{mN}$, $b_1, \dots, b_N \in \overline{U}$ we have $\sigma(b_1, \dots, b_N) = (b'_1, \dots, b'_N)$ where $(t - b_1) \dots (t - b_N) = (t - b'_1) \dots (t - b'_N)$ and $\overline{S}(b'_i) = \sigma \overline{S}(b_i)$, $\overline{S}$ stands for the ordered set of eigenvalues.

This action is local in the following sense. The transposition $(i, i + 1)$ for $\alpha m < i < (\alpha + 1)m$ acts only inside the $\alpha + 1$ -th factor of $\bar{U}^N$ and the transposition $(\alpha m, \alpha m + 1)$ acts only inside the product of the α -th and the $\alpha + 1$ -th factors. We have also the twisted transposition $\mu : \bar{U} \times \bar{U} \rightarrow \bar{U} \times \bar{U}$ in this case which is the action of the element $(1, m + 1)(2, m + 2) \dots (m - 1, 2m - 1) \in S_{2m}$.

Remark Let U be the set of matrix polynomials of the form $at + b$, where $a, b \in Mat_m$, $a = (a_{ij})$, $b = (b_{ij})$ and $a_{ij} = 0$ for $i < j$, $b_{ij} = 0$ for $i > j$. It is possible to define the twisted transposition μ such that for $\mu(f(t), g(t)) = (f_1(t), g_1(t))$ we have $f(t)g(t) = f_1(t)g_1(t)$, $\det f(t)$ and $\det g_1(t)$ have the same set of roots and the first coefficients of $f(t)$ and $g_1(t)$ have the same diagonal elements. In [4] we found the solutions of the corresponding twisted Yang-Baxter relation (for $m = 2$), which is generalization of the R -matrix from chiral Potts model.

§4. Local action of the symmetric group and factorization on matrix θ -functions

Let $\Gamma \subset \mathbb{C}$ be the lattice generated by 1 and τ where $\text{Im}\tau > 0$. We have $\Gamma = \{\alpha + \beta\tau; \alpha, \beta \in \mathbb{Z}\}$. Let $\varepsilon \in \mathbb{C}$ be primitive root of unity of degree m . Let $\gamma_1, \gamma_2 \in \text{Mat}_m$ be $m \times m$ matrices such that $\gamma_1^m = \gamma_2^m = 1, \gamma_2\gamma_1 = \varepsilon\gamma_1\gamma_2$. We have $\gamma_1 v_\alpha = \varepsilon^\alpha v_\alpha, \gamma_2 v_\alpha = v_{\alpha+1}$ in some basis $\{v_\alpha; \alpha \in \mathbb{Z}/m\mathbb{Z}\}$ of $\mathbb{C}^m$. Let us assume that $\{v_1, \dots, v_m\}$ is the standard basis of $\mathbb{C}^m$.

We denote by $M\Theta_{n,m,c}(\Gamma)$ for $n, m \in \mathbb{N}, c \in \mathbb{C}$ the space of everywhere holomorphic functions $f : \mathbb{C} \rightarrow \text{Mat}_m$, which satisfy the following equations:

$$\begin{aligned} f(z + \frac{1}{m}) &= \gamma_1^{-1} f(z) \gamma_1 \\ f(z + \frac{1}{m}\tau) &= e^{-2\pi i(mnz - c)} \gamma_2^{-1} f(z) \gamma_2 \end{aligned} \quad (6)$$

Proposition 3. $\dim M\Theta_{n,m,c}(\Gamma) = m^2 n$ and for each element $f \in M\Theta_{n,m,c}(\Gamma)$ the equation $\det f(z) = 0$ has exactly mn zeros modulo $\frac{1}{m}\Gamma$. The sum of these zeros is equal to $mc + \frac{mn}{2}$ modulo Γ .

Proof For $m = 1$ we have the usual θ -functions $\Theta_{n,c}(\Gamma) = M\Theta_{n,1,c}(\Gamma)$ and all these statements are well known in this case ([8]). One have a basis $\{\theta_\alpha(z); \alpha \in \mathbb{Z}/n\mathbb{Z}\}$ in the space $\Theta_{n,c}(\Gamma)$ such that $\theta_\alpha(z + \frac{1}{n}) = e^{2\pi i \frac{\alpha}{n}} \theta_\alpha(z), \theta_\alpha(z + \frac{1}{n}\tau) = e^{-2\pi i(z - \frac{n-1}{2n}\tau - \frac{1}{n}c)} \theta_{\alpha+1}(z)$ ([8]). From (6) it follows that $f(z + 1) = f(z)$ and $f(z + \tau) = e^{-2\pi i(m^2 nz - c_1)} f(z)$ for some $c_1 \in \mathbb{C}$. So the matrix elements of $f(z)$ are θ -functions from the space $\Theta_{m^2 n, c_1}(\Gamma)$. We have decomposition $f(z) = \sum_\alpha \varphi_\alpha \theta_\alpha(z)$, where $\varphi_\alpha \in \text{Mat}_m$ are constant matrices, $\{\theta_\alpha\}$ is a basis in the space $\Theta_{m^2 n, c_1}(\Gamma)$. Substituting this decomposition in (6) one can calculate the dimension of the space $M\Theta_{n,m,c}(\Gamma)$. We have also $\det f(z + \frac{1}{m}) = \det f(z)$ and $\det f(z + \frac{1}{m}\tau) = e^{-2\pi i(m^2 nz - mc)} \det f(z)$. From this follows the statement about zeros of the equation $\det f(z) = 0$.

Proposition 4. For generic complex numbers $\lambda_1, \dots, \lambda_{mn}$ such that $\lambda_1 + \dots + \lambda_{mn} \equiv mc + \frac{mn}{2} \pmod{\frac{1}{m}\Gamma}$ and nonzero vectors $v_1, \dots, v_{mn} \in \mathbb{C}^m$ there exist a unique up to proportionality element $f(z) \in M\Theta_{n,m,c}(\Gamma)$ such that $\det f(\lambda_\alpha) = 0, f(\lambda_\alpha)v_\alpha = 0$ for $1 \leq \alpha \leq mn$.

Proof Considering the decomposition $f(z) = \sum_\alpha \varphi_\alpha \theta_\alpha(z)$ one has the system of linear equations $\{\sum_\alpha \theta_\alpha(\lambda_\beta) \varphi_\alpha v_\beta = 0; \beta = 1, \dots, mn\}$ for matrix elements of $\{\varphi_\alpha\}$. One can see that this system defines $\{\varphi_\alpha\}$ uniquely up to proportionality for generic $\lambda_1, \dots, \lambda_{mn}, v_1, \dots, v_{mn}$.

We denote by $S(f)$ the set of zeros of the equation $\det f(z) = 0$ modulo $\frac{1}{m}\Gamma$.

Proposition 5. Let $f(z) \in M\Theta_{n,m,c}(\Gamma)$ be generic element and we have factorization $f(z) = f_1(z) \dots f_n(z)$ where $f_\alpha(z) \in M\Theta_{1,m,c_\alpha}(\Gamma)$, $c_1 + \dots + c_n = c$. Then $S(f_\alpha) \cap S(f_\beta) = \emptyset$ for $\alpha \neq \beta$ and $S(f) = S(f_1) \cup \dots \cup S(f_n)$. For each decomposition $S(f) = A_1 \cup \dots \cup A_n$ such that $A_\alpha \cap A_\beta = \emptyset$ for $\alpha \neq \beta$ and $\#A_\alpha = m$ there exists

a unique factorization $f(z) = f_1(z) \dots f_n(z)$ up to proportionality of f_α such that $S(f_\alpha) = A_\alpha$ for $\alpha = 1, \dots, n$.

Proof is similar to the proof of proposition 1, we just change polynomials by θ -functions.

Let U_c be the projectivisation of the linear space $M\Theta_{1,m,c}(\Gamma)$ and $U = \bigcup_{c \in \mathbb{C}} U_c$. We have the following twisted transposition $\mu : U \times U \rightarrow U \times U$. By definition $\mu(f, g) = (f_1, g_1)$, where $f(z)g(z) = f_1(z)g_1(z)$ and $S(f_1) = S(g), S(g_1) = S(f)$.

Let $\overline{U}$ be the set of elements f from U with fixed order on $S(f)$. For $f \in \overline{U}$ let $\overline{S}(f)$ be the set $S(f)$ with corresponding order. We have a local action of the symmetric group S_{mN} on the space $\overline{U}^N$. By definition, for $\sigma \in S_{mN}$ we have $\sigma(f_1, \dots, f_N) = (f_1^\sigma, \dots, f_N^\sigma)$ where $f_1(z) \dots f_N(z) = f_1^\sigma(z) \dots f_N^\sigma(z)$ and $\overline{S}(f_\alpha^\sigma) = \sigma \overline{S}(f_\alpha)$ for $1 \leq \alpha \leq N$.

Remark It is possible to construct twisted R -matrices for this twisted transposition μ as intertwiners of tensor products of cyclic representations of the algebra of monodromy matrices for elliptic Belavin R -matrix [5] at the point of finite order (see also [6]). It will be the subject of another paper.

§5. Generalization of the star-triangle relation

Let U be a complex manifold with an action of the symmetric group S_m by birational automorphisms. Let g_i be a birational automorphism of U corresponding to the transposition $(i, i+1)$, where $1 \leq i < m$. We assume also that the group S_{2m} acts on the manifold $U \times U$ by birational automorphisms, such that the transposition $(i, i+1)$ for $i < m$ acts by $g_i \times \text{id}$ and the transposition $(i, i+1)$ for $i > m$ acts by $\text{id} \times g_{i-m}$. Let f be the birational automorphism of $U \times U$ corresponding to the transposition $(m, m+1)$. We will use the notations $f(u, v) = (\lambda(u, v), \mu(u, v))$. We have: $g_i^2 = f^2 = \text{id}$, $g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1}$, $f g_m^l f = g_m^l f g_m^l$, $f g_1^r f = g_1^r f g_1^r$, where $g_m^l = g_m \times \text{id}$ and $g_1^r = \text{id} \times g_1$.

It is clear that we have the local action of the symmetric group S_{mN} on the manifold U^N for each N .

Let V be a n -dimensional vector space. For each $u \in U$ we denote by $V(u)$ a vector space canonically isomorphic to V . Let G_i ($1 \leq i < m$) be a meromorphic function from U to $\text{End}(V)$ and F be a meromorphic function from $U \times U$ to $\text{End}(V \otimes V)$. We will consider $G_i(u)$ and $F(u, v)$ as linear operators

$$G_i(u) : V(u) \rightarrow V(g_i(u))$$

$$F(u, v) : V(u) \otimes V(v) \rightarrow V(\lambda(u, v)) \otimes V(\mu(u, v))$$

Definition We call G_i, F a twisted GF-matrices if the following properties hold:

1. The compositions

$$V(u) \rightarrow V(g_i(u)) \rightarrow V(u)$$

$$V(u) \otimes V(v) \rightarrow V(\lambda(u, v)) \otimes V(\mu(u, v)) \rightarrow V(u) \otimes V(v)$$

are equal to the identity.

2. The following diagrams are commutative:

$$\begin{array}{ccc}
 V(g_i(u)) & \xrightarrow{G_{i+1}(g_i(u))} & V(g_{i+1}g_i(u)) \\
 \uparrow G_i(u) & & \downarrow G_i(g_{i+1}g_i(u)) \\
 V(u) & & V(g_i g_{i+1} g_i(u)) \\
 \downarrow G_{i+1}(u) & & \uparrow G_{i+1}(g_i g_{i+1}(u)) \\
 V(g_{i+1}(u)) & \xrightarrow{G_i(g_{i+1}(u))} & V(g_i g_{i+1}(u))
 \end{array}$$

$$\begin{array}{ccc}
V(\lambda(u, v)) \otimes V(\mu(u, v)) & \xrightarrow{G_m(\lambda(u, v)) \otimes 1} & V(g_m(\lambda(u, v))) \otimes V(\mu(u, v)) \\
\uparrow F(u, v) & & \downarrow F(g_m(\lambda(u, v)), \mu(u, v)) \\
V(u) \otimes V(v) & & V(\lambda(g_m(\lambda(u, v)), \mu(u, v))) \otimes V(\mu(g_m(\lambda(u, v)), \mu(u, v))) \\
\downarrow G_m(u) \otimes 1 & & \uparrow G_m(\lambda(g_m(u), v)) \otimes 1 \\
V(g_m(u)) \otimes V(v) & \xrightarrow{F(g_m(u), v)} & V(\lambda(g_m(u), v)) \otimes V(\mu(g_m(u), v))
\end{array}$$

$$\begin{array}{ccc}
V(\lambda(u, v)) \otimes V(\mu(u, v)) & \xrightarrow{1 \otimes G_1(\mu(u, v))} & V(\lambda(u, v)) \otimes V(g_1(\mu(u, v))) \\
\uparrow F(u, v) & & \downarrow F(\lambda(u, v), g_1(\mu(u, v))) \\
V(u) \otimes V(v) & & V(\lambda(\lambda(u, v), g_1(\mu(u, v)))) \otimes V(\mu(\lambda(u, v), g_1(\mu(u, v)))) \\
\downarrow 1 \otimes G_1(v) & & \uparrow 1 \otimes G_1(\mu(u, g_1(v))) \\
V(u) \otimes V(g_1(v)) & \xrightarrow{F(u, g_1(v))} & V(\lambda(u, g_1(v))) \otimes V(\mu(u, g_1(v)))
\end{array}$$

It is clear that we have also twisted transposition μ in this situation. By definition μ is the birational automorphism of $U \times U$ corresponding to the element $(1, m+1)(2, m+2) \dots (m-1, 2m-1) \in S_{2m}$. We have

$$\mu = \prod_{1 \leq \alpha \leq m} g_{m-\alpha+1}^l \dots g_{m-1}^l f g_{m-1}^r \dots g_\alpha^r$$

where $g_\alpha^l = g_\alpha \times \text{id}$, $g_\alpha^r = \text{id} \times g_\alpha$ and the product is ordered from $\alpha = 1$ to $\alpha = m$. One can also construct the corresponding twisted R -matrix as a product of m^2 factors:

$$R = \prod_{1 \leq \alpha \leq m} (G_{m-\alpha+1} \otimes 1) \dots (G_{m-1} \otimes 1) F(1 \otimes G_{m-1}) \dots (1 \otimes G_\alpha) \quad (7)$$

Here we omit arguments of R, G_α and F .

Let W be a vector space, L be a meromorphic function from U to $\text{End}(V \otimes W)$. We will consider $L(u)$ as a linear operator $L(u) : V(u) \otimes W \rightarrow W \otimes V(u)$.

Definition We call L an L -operator for twisted GF -matrices G_i, F if the following diagrams are commutative:

$$\begin{array}{ccc}
V(u) \otimes W & \xrightarrow{L(u)} & W \otimes V(u) \\
\downarrow G_i(u) \otimes 1 & & \downarrow 1 \otimes G_i(u) \\
V(g_i(u)) \otimes W & \xrightarrow{L(g_i(u))} & W \otimes V(g_i(u))
\end{array}$$

$$\begin{array}{ccc}
V(u) \otimes W \otimes V(v) & \xrightarrow{L(u) \otimes 1} & W \otimes V(u) \otimes V(v) \\
\uparrow 1 \otimes L(v) & & \downarrow 1 \otimes F(u, v) \\
V(u) \otimes V(v) \otimes W & & W \otimes V(\lambda(u, v)) \otimes V(\mu(u, v)) \\
\downarrow F(u, v) \otimes 1 & & \uparrow L(\lambda(u, v)) \otimes 1 \\
V(\lambda(u, v)) \otimes V(\mu(u, v)) \otimes W & \xrightarrow{1 \otimes L(\mu(u, v))} & V(\lambda(u, v)) \otimes W \otimes V(\mu(u, v))
\end{array}$$

That is

$$\begin{aligned}
G_i(u)L(u) &= L(g_i(u))G_i(u), \text{ for } 1 \leq i < m \\
F(u, v)L^1(u)L^2(v) &= L^1(\lambda(u, v))L^2(\mu(u, v))F(u, v)
\end{aligned} \tag{8}$$

We can consider $L(u)$ as a $n \times n$ matrix $L(u) = (L_i^j(u))$, where $L_i^j(u)$ are the elements in $End(W)$ such that

$$L(u) : x_i(u) \otimes W \rightarrow (L_i^j(u)W) \otimes x_j(u)$$

Here $\{x_i\}$ is a basis in V and $\{x_i(u)\}$ is the corresponding basis in $V(u)$.

Definition The algebra of monodromy matrices M_{GF} for the twisted GF -matrices G_i, F is an associative algebra with the generators $\{L_i^j(u); i, j = 1 \dots n, u \in U\}$ and the defining relations (8).

It is clear that the representations of the algebra M_{GF} in the space W coincide with the L -operators $L(u) \in End(V \otimes W)$.

Since the relations (4) follow from the relations (8), one has a homomorphism $M_{GF} \rightarrow M_R$, which is identity on the generators, where R is given by (7).

It is clear that the algebra M_{GF} has the usual coproduct $\Delta : L_i^j(u) \rightarrow L_i^\alpha(u) \otimes L_\alpha^j(u)$.

Remarks 1. One can also construct the Zamolodchikov algebra for GF -matrices.

2. For $Q(u) = \text{tr} L(u) = L_i^i(u)$ we have the following relations:

$$\begin{aligned}
Q(u) &= Q(g_i(u)) \text{ for } 1 \leq i < m \\
Q(u)Q(v) &= Q(\lambda(u, v))Q(\mu(u, v))
\end{aligned}$$

3. It is possible to construct twisted GF -matrices for the elliptic local action of the symmetric group defined in §4 as intertwiners of tensor products of cyclic representations of the algebra of monodromy matrices for elliptic Belavin R -matrix ([5]) at the point of finite order. It will be the subject of another paper.

Acknowledgments

I am grateful to V.Bazhanov and A.Belavin for useful discussions.

I am grateful to Max-Planck-Institut für Mathematik, Bonn where this paper was written for invitation and very stimulating working atmosphere.

The work is supported partially by RFBR 99-01-01169, RFBR 00-15-96579, CRDF RP1-2254 and INTAS-00-00055.

References

1. R.J. Baxter, J.H.H. Perk and H. Au-Yang, New solutions of the star-triangle relations for the chiral Potts model, *Phys. Lett. A*, 128(1988)138-142.
2. V.V. Bazhanov and Yu.G. Stroganov, Chiral Potts model as a descendant of the six vertex model, *J. Stat. Phys.* 51(1990)799-817.
3. V.O. Tarasov, Cyclic monodromy matrices for the R -matrix of the six-vertex model and the chiral Potts model with fixed spin boundary conditions, *International Journal of Modern Physics A*, Vol.7, Suppl. 1B(1992)963-975.
4. A.A. Belavin, A.V. Odesskii and R.A. Usmanov, New relations in the algebra of the Baxter Q -operators. hep-th/0110126.
5. A.A. Belavin, Discrete groups and integrability of quantum systems, *Func. Anal. Appl.* 14(1980)18-26.
6. B.L. Feigin and A.V. Odesskii, Sklyanin Elliptic Algebras. The case of point of finite order, *Func. Anal. Appl.* 29(1995).
7. I. Gelfand, S. Gelfand, V. Retakh, S. Serconek, R.L. Wilson, Hilbert series of quadratic algebras associated with pseudo-roots of noncommutative polynomials, QA/0109007.
8. Mumford, Tata lectures on theta. I. With the assistance of C. Musili, M. Nory and M. Stillman. *Progress in Mathematics*, 28. BirkhDuser Boston, Inc., Boston, MA, 1983.